

DERWENT PUBLICATIONS LTD.

46109

<p>46109 E/23 HUMINAL VERTIEBSGE 23.10.80-DE-040040 (03.06.82) C05f-11 C05g-01 C09k-17 Fertiliser prep. by composting conifer bark - by mixing ground bark with ammonium bicarbonate and water</p>	<p>C(4-A7D, 5-C1, 12-N8, 12-N9, 12-N10) D(5-A4) : 3</p> <p>with the addn. of plant nutrients, to promote plant growth. Pref. the end prod. has total N : P₂O₅ : K₂O (insol) of 1 : 1 : 1.5 or 2 : 1 : 1.5 or 4 : 1 : 1.5.</p>
<p>Prepn. of a soil improving agent by composting conifer bark, comprises grinding fresh conifer bark to particle size 0-18 mm. and then mixing with 1-2 kg. (per 1000 l. fresh bark) of pure nitrogen in the form of ammonium bicarbonate, and with 70-80 wt. % water to give a homogeneous mixt. which is then composted.</p> <p><u>USES/ADVANTAGES</u> The fertiliser smells like fresh forest soil, is dark in colour and is a good absorber of water (62% at pH 0.6). None of these properties are possessed by calcium nitrate as nitrogen carrier. The bacteria present in the bark need nitrogen to synthesise their protein; ammonia is a considerably better source of this than nitrate.</p> <p><u>DETAILS</u> The conifer bark is pref. from a spruce, fir tree or pine tree. Pref. the composted bark is enriched with a 25-50 vol. % peat with degree of decomposition H6 to H10, opt.</p>	<p><u>EXAMPLE</u> Bark was mixed with 6 kg. ammonium bicarbonate (17.5 % N) per 1000 l. fresh bark and the water content set to 70% at 20°C. After 1 week the temp. rose to 65°C. and reaction continued until the 5th week; when the temp. reached 70°C., after which the temp. fell, until in the 12th week the temp. had stabilised.(18pp1401).</p>

DE3040040

BEST AVAILABLE COPY